

## Identification of Drought Tolerant Maize Genotypes and Seedling based Morpho-Physiological Selection Indices for Crop Improvement

(Pengenalpastian Genotip Jagung yang Tahan Kemarau dan Indeks Pemilihan Morfo-Fisiologi berasaskan Anak Benih untuk Pembaikan Tanaman)

FAHAD MASOUD WATTOO, RASHID MEHMOOD RANA, SAJID FIAZ\*, SYED ADEEL ZAFAR, MEHMOOD ALI NOOR, HAFIZ MUMTAZ HASSAN, MUHAMMAD HUSNAIN BHATTI, SHOAB UR REHMAN, GALAL BAKR ANIS & RAI MUHAMMAD AMIR

### ABSTRACT

*Maize is an imperative grain crop used as a staple food in several countries around the world. Water deficiency is a serious problem limiting its growing area and production. Identification of drought tolerant maize germplasm is comparatively easy and sustainable approach to combat this issue. Present research was conducted to evaluate 50 maize genotypes for drought tolerance at early growth stage. Drought tolerance was assessed on the basis of several morphological and physiological parameters. Analysis of variance showed significant variation among the tested maize genotypes for recorded parameters. Principal component analysis revealed important morpho-physiological traits that were playing key role in drought tolerance. Correlation studies depicted significant positive correlation among the attributes such as fresh shoot length (FSL), fresh root length (FRL), dry shoot weight (DSW), dry root weight (DRW), relative water contents (RWC) and total dry matter (TDM) while a strongly negative correlation was observed among RWC and excised leaf water loss. Results concluded that the parameters fresh shoot weight, fresh root weight, FRL, DRW, TDM, cell membrane thermo stability (CMT) and RWC can be useful for rapid screening of maize germplasm for drought tolerance at early growth stages. Furthermore, the genotypes 6, 16, 18, 40, 45 and 50 can be used as a drought tolerant check in breeding programs. Moreover, biplot analysis along with other indices was proved to be a useful approach for rapid and cost efficient screening of large number of genotypes against drought stress condition.*

*Keywords: Cell membrane thermo stability; correlation; drought tolerance; principal component analysis; relative water contents*

### ABSTRAK

*Jagung adalah tanaman bijirin penting yang digunakan sebagai makanan ruji di beberapa negara di seluruh dunia. Kekurangan air adalah masalah serius yang membatasi kawasan dan pengeluarannya yang semakin meningkat. Pengenalpastian germplasma jagung yang tahan kemarau adalah pendekatan yang agak mudah dan mampan untuk memerangi isu ini. Penyelidikan ini dijalankan untuk menilai 50 genotip jagung untuk toleransi kemarau pada peringkat pertumbuhan awal. Ketahanan kemarau dinilai berdasarkan beberapa parameter morfologi dan fisiologi. Analisis varians menunjukkan variasi ketara antara genotip jagung yang diuji untuk parameter yang direkodkan. Analisis komponen utama mendedahkan ciri morfo-fisiologi yang memainkan peranan penting dalam ketahanan kemarau. Kajian korelasi menunjukkan korelasi positif yang signifikan antara ciri seperti panjang pucuk segar (FSL), panjang akar segar (FRL), berat pucuk kering (DSW), berat akar kering (DRW), kandungan air relatif (RWC) dan jumlah bahan kering (TDM) manakala korelasi yang sangat negatif diperhatikan dalam kalangan RWC dan mengurangkan kehilangan air daun. Keputusannya menyimpulkan bahawa parameter pucuk berat baru, berat akar segar, FRL, DRW, TDM, kestabilan thermo membran sel (CMT) dan RWC adalah berguna untuk penapisan pantas germplasma jagung untuk ketahanan kemarau pada peringkat pertumbuhan awal. Selain itu, genotip 6, 16, 18, 40, 45 dan 50 boleh digunakan sebagai pemeriksaan ketahanan kemarau dalam program pembiakan. Selain itu, analisis biplot bersama-sama dengan indeks lain telah terbukti merupakan pendekatan yang berguna untuk penapisan pantas dan cekap kos untuk sejumlah besar genotip terhadap keadaan tekanan kemarau.*

*Kata kunci: Analisis komponen utama; kandungan air relatif; kestabilan thermo membran sel; ketahanan kemarau; korelasi*

### INTRODUCTION

Maize is an imperative cereal crop of the world serving as a source of food and feed for humans and animals,

respectively. Several biotic and abiotic stresses affect its growth and yield (Abbasi et al. 2016; Waqas et al. 2017; Zafar et al. 2015). However, drought stress has been

the major limiting factor affecting its yield potential (Khodarahmpour & Hamidi 2011; Seghatoleslami et al. 2008). It affects various morpho-physiological processes including plant biomass, root length, shoot length, photosynthesis, water use efficiency (WUE) and leaf water content (Egilla et al. 2005; Jaleel et al. 2008).

Screening for drought tolerant germplasm is a pre-requisite for any breeding program. However, assessment of germplasm for abiotic stresses like heat and drought under natural field conditions is difficult due to uncertain environmental conditions like rainfall and humidity. (Zafar et al. 2017). The task becomes more challenging when dealing with large germplasm due to requirement of large area, labor, resources and long time period. Alternatively, its rapid evaluation can be done at early growth stages under controlled environmental conditions based on several growth and physiological parameters (Zafar et al. 2017). Various growth related to morpho-physiological traits have been potentially utilized for evaluating genotypes of different crops against water stress (Ali et al. 2009; Chohan et al. 2012; Javed 2012; Taiz & Zeiger 2006). Root to shoot ratio of plants has been reported to increase under water deficit conditions (Wu & Cosgrove 2000). This ratio deviates because roots are less sensitive to growth inhibition by low water potential as compared to shoots. Significant variation exists in maize seedlings for root to shoot ratio and shoot dry weight under water deficit condition (Gonzales & Ayerbe 2011). Dry root weight was found to be a valuable selection criterion for selection against water stress (Mehdi et al. 2001). Assessment of drought tolerance in cereals indicated a significant positive correlation for root length with grain weight, grain yield (Shrimali 2001) and harvest index (Kashiwagi et al. 2005). Among various water stress resistance traits targeted in crop breeding, cell membrane thermo-stability (CMT) (Chohan et al. 2012) and coleoptile length (Qayyum et al. 2011) have been considered as the key physio-morphological traits affected under drought (Bayoumi et al. 2008).

Selection for best performing genotypes under stress condition is the primary objective for any crop improvement program (Thiry et al. 2016). However, low heritability of drought tolerant traits and unavailability of effective selection approach limits the development of drought tolerant cultivars (Kirigwi et al. 2004). Various morphological, physiological and biological features of higher plants show adaptability in response to water stress. Existence of variability for drought tolerance in different plant species has been reported (Frova et al. 1999; Gutteri et al. 2004). According to 'stress gradient hypothesis' the results of seedling analysis determine the structure and dynamics of plant population (De La Cruz et al. 2008; Kitajima & Fenner 2000). Therefore, phenotypic evaluation at the seedling stage is regarded an attractive approach having high throughput and cost efficient, saves space and time (Meeks et al. 2013). This approach has been successfully utilized to develop drought tolerant varieties in different crop species (Meeks et al. 2013).

When dealing with a large number of genotypes, multivariate biometrical techniques are utilized to assess biodiversity irrespective of data set. Among these biometrical techniques, biplot analysis or principal component analysis (PCA), principal coordinate analysis (PCoA), cluster analysis and multidimensional scaling (MDS) are utilized by plant breeders (Brown-Guedira et al. 2000). Biplot analysis interpreted easily and requires only a glance to understand the tolerance potential of genotypes due to graphical presentation of data. Moreover, biplot also illustrates average or overall performance of genotypes across the environments which makes it a reliable technique (Ali et al. 2012).

Present study was conducted with the objectives: to identify the genetic variation in maize germplasm and genotypes which can perform well under limited water conditions, and to evaluate the various morpho-physiological parameters related to drought tolerance which can be used efficiently as selection criteria in breeding programs.

#### MATERIALS AND METHODS

A set of 50 diverse maize inbred lines were collected from various national and international research organizations (Table 1). The genotypes were evaluated for water stress tolerance on the basis of various seedling based morpho-physiological parameters. The seeds were sown in plastic pots of one liter (L) volume filled with sand during spring 2014 in completely randomized design (CRD) in wire house with three replications under both normal and water deficit conditions. One seedling in each pot was maintained through thinning after uniform emergence. Five seedlings of each genotype were kept in each replication and equal amount of water was applied to facilitate germination to each pot in both the treatments. The first irrigation was applied after seven days of sowing to both normal and water deficit treatments. After two weeks of sowing the second irrigation was supplied to entries only under normal condition. Seedlings under both treatments were allowed to grow further for seven days. Data pertaining to physiological and morphological parameters were recorded before and after uprooting seedlings by applying adequate amount of water to entries in both treatments. The average range for minimum and maximum temperature throughout experimental duration in the wire house was  $30 \pm 2$  °C and  $35 \pm 2$  °C, respectively.

Data were recorded for various traits before uprooting of seedlings. Coleoptile length was recorded after seven days of sowing and was measured from the base to the tip of coleoptile in evening about 18:00 h in centimeters (cm). Leaf temperature (LT) was measured for each treatment at 13:00 - 15:00 h from fully expanded leaves. Measurements were recorded from three leaves of each seedling with infrared thermometer (RAYPRM 30 CFRJ, RAYTEK, USA).

CMT an indicator of relative cell injury percentage was determined according to protocol described elsewhere

(Ibrahim & Quick 2001). Relative water contents (RWC) were determined from five leaves of each sample per treatment using the standard method (García-Mata & Lamattina 2001). Fresh, dry and turgid weight from leaves were measured by using standard methods. Relative water contents were determined by using the formula.

$$\text{RWC \%} = \frac{[\text{Fresh Leaf Weight} - \text{Dry Leaf Weight}]}{[\text{Turgid Leaf Weight} - \text{Dry Leaf Weight}]} \times 100.$$

Excised leaf weight loss (ELWL) were determined by using following formula (Clarke & Townley-Smith 1986).

$$\text{ELWL} = \frac{(\text{Fresh Leaf Weight} - \text{weight after 6 h})}{(\text{Fresh Leaf Weight} - \text{Dry Leaf weight})} \times 100.$$

Fresh root weight, fresh shoot weight, fresh root length and fresh shoot length were recorded immediately after uprooting the seedling. Dry root weight, dry shoot weight and total dry matter were recorded after complete drying of samples in incubator. Root to shoot ratio (RSR) was obtained according to the formula described elsewhere (Nour & Weibel 1978).

Root to shoot ratio = Dry root weight/Dry shoot weight

Genotypic and phenotypic variance and heritability were calculated by using following formulas.

$$\text{Genotypic variance } (\sigma^2g) = \frac{[\text{Genotype mean square (GMS)} - \text{Error mean square (EMS)}]}{\text{number of replications (r)}}$$

$$\text{Phenotypic variance } (\sigma^2p) = \sigma^2g + \sigma^2e/r$$

$$\text{Heritability } (H^2) = \sigma^2g/\sigma^2p.$$

#### STATISTICAL ANALYSIS

Data related to seedling based morphological and physiological traits were analyzed using GenStat version 10 for analysis of variance (ANOVA) and GGE biplot analysis. XL-stat version 2012 software was used to compute the correlation analysis.

#### RESULTS AND DISCUSSION

Analysis of variance (ANOVA) showed significant variation among the tested maize genotypes for recorded parameters

TABLE 1. List of Maize germplasm under study evaluated for drought tolerance

Sr. No.	Inbred lines	Sr. No.	Inbred lines
1	NCIL-10-4	26	D-130
2	NCIL-10-5	27	D-135
3	NCIL-10-23	28	D-132
4	NCIL-20-4	29	D-133
5	NCIL-20-11	30	D-134
6	NCIL-20-20	31	D-136
7	NCIL-30-4	32	D-137
8	NCIL-30-5	33	D-143
9	NCIL-30-15	34	D-148
10	NCIL-40-8	35	D-149
11	NCIL-40-12	36	D-150
12	NCIL-40-18	37	D-152
13	D-103	38	D-153
14	D-105	39	D-155
15	D-106	40	D-157
16	D-109	41	D-158
17	D-111	42	W 64 TMS
18	D-114	43	W 64 SP
19	D-118	44	OH 54-3A
20	D-120	45	OH-8
21	D-122	46	Antigua-1
22	D-123	47	AES-204
23	D-126	48	A-638
24	D-131	49	A-556
25	D-124	50	M-14

TABLE 2. Mean Square of various morpho physiological traits in 50 Maize genotypes evaluated for drought tolerance

	MS	$\sigma^2e$	$\sigma^2g$	$\sigma^2p$	ECV%	GCV%	PCV%	H2%	GA
RSRW	1.88**	0.083	0.9	0.98	17.601	57.72	60.34	91.49	1.86
TDM	7.14**	0.074	3.53	3.61	6.72	46.27	46.76	97.93	3.83
CL	6.227**	0.55	2.83	3.39	19.82	44.86	49.05	83.67	3.17
CMT	243.47**	21.45	111.09	132.464	11.7	26.63	29.1	83.8	19.86
RWC	246.73**	5.49	120.61	126.11	3.56	16.68	17.06	95.64	28.57
ELWL	324.44*	24.77	149.83	174.61	12.29	30.23	32.64	85.81	23.35
LT	9.04*	2.8	3.11	5.92	4.57	4.82	6.65	52.65	2.63
FSL	25.62*	0.769	12.42	13.19	3.12	12.54	12.92	94.17	9.09
FRL	43.708**	0.76	21.47	22.23	3.19	16.99	17.29	96.58	9.38
FSW	29.56**	0.243	14.62	14.9	5.11	39.6	39.93	98.36	7.822
FRW	23.65**	0.58	11.53	12.11	8.31	36.96	37.89	95.18	6.82
DSW	0.180*	0.058	0.061	0.119	15.732	16.099	22.509	51.15	0.47
DRW	6.154**	0.027	3.063	3.091	6.61	69.27	69.58	99.1	3.58

\* = significance at 0.05 while \*\* = significant at 0.01 percent probability level, respectively. MS = Mean square  $\sigma^2e$  = environmental variance,  $\sigma^2g$  = genotypic variance  $\sigma^2p$  = phenotypic variance, H2= broad sense heritability, GA= genetic advance, RSR = root shoot ratio, TDM= total dry matter, CL= coleoptile length, CMT= cell membrane thermo-stability, RWC= relative water content, ELWL= excised leaf weight loss, LT= leaf temperature, FSL= fresh shoot length, FRL= fresh root length, FSW= fresh shoot weight, FRW= fresh root weight, DSW= dry shoot weight, DRW= dry root weight

(Table 2), pointing out considerable genetic diversity in germplasm.

#### COEFFICIENT OF VARIATION AND VARIANCE COMPONENTS

The estimates of genetic traits such as genotypic mean square (MS), broad sense heritability ( $H^2$ ), genetic advance (GA), genotypic, phenotypic and environmental variances ( $\sigma^2g$ ,  $\sigma^2p$  and  $\sigma^2e$ , respectively) along with genotypic, phenotypic and environmental coefficient of variation (GCV, PCV and ECV, respectively) are given in Table 2. For all the traits under investigation, values of phenotypic variance were greater as compared to genotypic variance. ELWL showed highest phenotypic and genotypic variance whereas, dry shoot weight (DSW) showed minimum genotypic and phenotypic variance. Similarly, values of phenotypic and genotypic variances were high as compared to environmental variance for all the recorded traits.

The outcome of present investigation showed that environmental variation was greater enough to modify expression of gene systematizing the trait under examination. Selection pressure might be applied on traits to spot superlative genotypes from given germplasm. Similar outcomes were obtained for GCV, PCV and ECV (Table 2). The differences between GCV and PCV were very less for almost all the traits under study which showed a small effect of environment on these traits suggesting that selection of these traits could be effective. Our results were compatible with previous findings (Ali et al. 2013; Mustafa et al. 2013). In current study, dry root weight (DRW) showed maximum PCV and GCV followed by RSR with PCV and GCV. These attributes can be used as a reliable selection criteria for drought tolerance in maize at early growth stage. On the other hand, LT showed minimum PCV and GCV followed by FSL with PCV and GCV. Thus, these parameters which showed minimum PCV and GCV were regarded as poor index for selection of genotypes against

drought tolerance. Our results were in line with a previous report (Ali et al. 2011).

#### HERITABILITY AND GENETIC ADVANCE OF TESTED MORPHO-PHYSIOLOGICAL PARAMETERS

Heritability is a good indicator for transmission of attributes from parents to offspring and scope of trait improvement through selection breeding relies on the value of heritability (Neelima & Reddy 2008). GCV coupled with heritability estimate presents a more reliable approximate of quantity of GA to be anticipated through phenotypic selection. Heritability values <59% are considered as low while values ranged between 60-79% are referred to as moderately higher. Whereas, values greater than 80% are considered as higher values (Singh & Singh 1999).

Higher heritability was observed for DRW, FSW, TDM, FRL, RWC and CMT, respectively (Table 2). Similar results were obtained in some previous investigations for these traits (Chohan et al. 2012; Rezaeieh & Eivazi 2011). RWC, ELWL, CMT and FRL showed higher GA. However, GA was less for DSW and RSR. Our results were in agreement with past findings (Mustafa et al. 2013; Nzube et al. 2014). The attributes under observation showed higher values for broad sense heritability and GA indicating additive gene action with less environmental influence. Thus, these traits can be employed as useful selection criteria for maize improvement programs.


#### PRINCIPAL COMPONENT ANALYSIS (PCA) REVEALED USEFUL PARAMETERS FOR DROUGHT TOLERANCE

Out of 13 principal components (PCs), first three PCs exhibited Eigen values more than 1 with a combined variance of 66.31% (Figure 1(A)) whereas, remaining 10 PCs showed only 33.68% cumulative variability. First two PCs showed a combined variance of 58.50% suggesting that these were playing key role in overall

variation. The PC 1 had a high match with DRW followed by TDM. While PC 2 was more related to FRL followed by FSW (Figure 1(B)). Factor loadings for genotypes showed that genotypes 50, 18, 16, 40, 6 and 45 exhibited higher values for PC 1 (Figure 2). Whereas, second PC was more related to genotype 42 followed by genotype 3 (Figure 2). Genotypes with higher PC 1 and lower PC 2 values are considered stable genotypes while genotypes with lower PC 1 and higher PC 2 are perceived as unstable genotypes (Seghatoleslami et al. 2008). Use of this biometrical technique for pin pointing water deficit stress tolerance maize genotypes has previously been reported in maize crop (Ahmadzadeh 1997). Since FRW, DRW, TDM, RWC, CMT, CL and RSR showed higher values in PC 1 and lower values in PC 2, so it is presumable that these traits can be used as a selection marker for drought tolerance in maize. Similarly, genotypes 50, 18, 16, 40, 6 and 45 showed higher values for first PC accompanied with lower second PC values; hence these genotypes can be considered as stable for selection breeding.

BIPLOT ANALYSIS SHOWED SIGNIFICANT DIVERSITY AMONG TESTED MAIZE GERMPLASM

In present investigation, biplot analysis revealed high genetic diversity among the tested maize genotypes for all



the recorded parameters (Figure 3). Closest the distance of genotype clog position on graph from the point of origin least will be the diversity and vice versa. Maize genotypes which are located at the vertex of the polygon were regarded as most diversified compared to others. Genotypes 50, 18, 45, 6, 11, 44, 14 and 42 clogged at the vertex of polygon hence considered more diversified and of more breeding value. Interestingly the genotypes 50, 18, 6, 16, 40 and 45 were also proved stable based on high PC1 and lower PC2 (Figure 2), thus can be utilized in maize improvement programs for drought tolerance. Moreover, the traits FRL, FSW, TDM, DSW, DRW and RSR were clogged away from the point of origin which depicted them as stable and reliable for selection against drought tolerance in maize.

CORRELATION ANALYSIS AMONG MORPHO-PHYSIOLOGICAL TRAITS FOR DROUGHT TOLERANCE

Correlation analysis revealed significant positive as well as negative association among several traits under examination (Table 3). A significant ( $p < 0.01$ ) positive correlation was observed between LT and ELWL. Previously, strong positive correlation was also observed between LT and ELWL (Khan et al. 2014). However, LT was negatively correlated with CMT, FSW, FRW, DSW and DRW.




FIGURE 1. Data showing eigen value more than 1 and cumulative percent variance of 66.31% in first three PCs (A), and factor loadings for traits (B) based on 50 maize genotypes evaluated under normal and water deficit conditions




FIGURE 2. Factor loadings for genotypes showing higher value in genotypes 50, 18, 16, 40, 6 and 45

TABLE 3. Correlation among various seedlings based morpho-physiological parameters under normal and water deficit conditions in maize

	RSRW	TDM	CL	CMT	RWC	ELWL	LT	FSL	FRL	FSW	FRW	DSW	DRW
RSRW	1												
TDM	0.88542	1											
CL	0.70801	0.78341	1										
CMT	0.36841	0.49074	0.50394	1									
RWC	0.44615	0.4844	0.56538	0.28333	1								
ELWL	-0.6576	-0.7812	-0.6297	-0.5113	-0.3815*	1							
LT	-0.4123	-0.3901	-0.3881	-0.1695	-0.2036	0.42216**	1						
FSL	0.46617*	0.45903	0.48992	0.23006	0.29303	-0.4307	-0.1993	1					
FRL	0.27359	0.22894	0.35427	0.08357	0.17863	-0.2624	-0.3709	0.37591*	1				
FSW	0.15399	0.12216	0.18813	0.23968	0.23191	-0.1039	-0.1696	0.13475	0.24696	1			
FRW	0.6191	0.65063	0.58797	0.41771	0.36433	-0.5988	-0.494	0.45575	0.2848	0.166	1		
DSW	0.08188 <sup>NS</sup>	0.51552	0.30144	0.26883	0.14176	-0.4114	-0.1185	0.07636	0.00456	-0.0904	0.21502	1	
DRW	0.94003	0.98915	0.79247	0.48271	0.49765	-0.7713	-0.4*	0.48153**	0.24591	0.14712	0.66421	0.38406*	1

\* = significance at 0.05 while \*\* = significant at 0.01 percent probability level, respectively. RSR = root shoot ratio, TDM= total dry matter, CL= coleoptile length, CMT= cell membrane thermo-stability, RWC= relative water content, ELWL= excised leaf weight loss, LT= leaf temperature, FSL= fresh shoot length, FRL= fresh root length, FSW= fresh shoot weight, FRW= fresh root weight, DSW= dry shoot weight, DRW= dry root weight




FIGURE 3. Biplot Analysis showing different attitude of 50 Maize genotypes for various morphological attributes under drought stress condition. \* = significance at 0.05 while \*\* = significant at 0.01 percent probability level, respectively

The reason for the increased ELWL under high LT seems to be the increased transpiration and reduced CMT level due to heat shock. FSL also showed significant positive association with FRL and DRW. Mustafa et al. (2013) also reported a significant positive association among FSL, FRL and DRW. Positive correlation was also noticed between DSW and DRW. Some previous studies were found

to support our results (Ali et al. 2011). There exists a negative correlation among RWC and ELWL (Amini et al. 2013). Present study also showed a significant negative correlation between RWC and ELWL. The reason for the decreased RWC under increased ELWL seems to be due to excessive water loss by high transpiration rate and reduced CMT.

## CONCLUSION

Significant variation was observed among the tested maize genotypes for drought tolerance using biplot analysis. Genotypes 6, 16, 18, 40, 45 and 50 were proved to be relatively drought tolerant based on tested morpho-physiological indices. In addition, FRL, FSW, FRW, DRW, TDM, CMT and RWC were identified as useful morpho-physiological indices based on principal component analysis along with higher heritability and GA. Moreover, Biplot analysis along with other indices was proved to be a useful approach for rapid and cost efficient screening of large number of genotypes against drought stress condition.

## REFERENCES

- Abbasi, G.H., Ijaz, M., Akhtar, J., Anwar Ul-Haq, M., Jamil, M., Ali, S., Ahmad, R. & Khan, H.N. 2016. Profiling of anti-oxidative enzymes and lipid peroxidation in leaves of salt tolerant and salt sensitive maize hybrids under NaCl and Cd stress. *Sains Malaysiana* 45(2): 177-184.
- Ahmazadeh, A. 1997. Determination of the best drought tolerance index in selected maize (*Zea mays* L.) lines, MSc. Thesis, Tehran University, Tehran, Iran (Unpublished).
- Ali, M.A., Niaz, S., Abbas, A., Sabir, W. & Jabran, K. 2009. Genetic diversity and assessment of drought tolerant sorghum landraces based on morpho-physiological traits at different growth stages. *Plant Omics* 2(5): 214-227.
- Ali, Q., Ahsan, M., Mustafa, H.S.B. & Hasan, E.U. 2013. Studies of genetic variability and correlation among morphological traits of maize (*Zea mays* L.) at seedling stage. *Albanian Journal of Agricultural Sciences* 12(3): 405-410.
- Ali, Q., Elahi, M., Ahsan, M., Tahir, M.H.N. & Basra, S.M.A. 2011. Genetic evaluation of maize (*Zea mays* L.) genotypes at seedling stage under moisture stress. *International Journal for Agro Veterinary and Medical Sciences* 5(2): 184-193.
- Ali, Z., Salam, A., Azhar, F.M., Khan, I.A., Khan, A.A., Bahadur, S., Mahmood, T., Ahmad, A. & Trethowan, R. 2012. The response of genetically distinct bread wheat genotypes to salinity stress. *Plant Breeding* 131(6): 707-715.
- Amini, Z., Khodambashi, M. & Houshmand, S. 2013. Correlation and path coefficient analysis of seed yield related traits in maize. *International Journal of Agriculture and Crop Sciences* 5(19): 2217-2220.
- Gonzales, A. & Ayerbe, L. 2011. Response of coleoptiles to water deficit: Growth, turgor maintenance and osmotic adjustment in barley plants (*Hordeum vulgare* L.). *Agricultural Sciences* 2(3): 159-166.
- Bayoumi, T.Y., Eid, M.H. & Metwali, E.M. 2008. Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes. *African Journal of Biotechnology* 7(14): 2341-2352.
- Brown-Guedira, G.L., Thompson, J.A., Nelson, R.L. & Warburton, M.L. 2000. Evaluation of genetic diversity of soybean introductions and North American ancestors using RAPD and SSR markers. *Crop Science* 40(3): 815-823.
- Chohan, M.S.M., Saleem, M., Ahsan, M. & Asghar, M. 2012. Genetic analysis of water stress tolerance and various morpho-physiological traits in (*Zea mays* L.) using graphical approach. *Pakistan Journal of Nutrition* 11(5): 489-500.
- Clarke, J.M. & Townley-Smith, T.F. 1986. Heritability and relationship to yield of excised-leaf water retention in durum wheat. *Crop Science* 26(2): 289-292.
- De La Cruz, M., Roberto, L.R., Adrián, E. & Maestre, F.T. 2008. Where do seedlings go? A spatio-temporal analysis of seedling mortality in a semi-arid gypsophyte. *Ecography* 31(6): 720-730.
- Egilla, J.N., Davies, F.T. & Boutton, T.W. 2005. Drought stress influences leaf water content, photosynthesis, and water-use efficiency of *Hibiscus rosa-sinensis* at three potassium concentrations. *Photosynthetica* 43(1): 135-140.
- Frova, C., Krajewski, P., Fonzo, N.D., Villa, M. & Sari-Gorla, M. 1999. Genetic analysis of drought tolerance in maize by molecular markers I. Yield components. *Theoretical and Applied Genetics* 99(1-2): 280-288.
- García-Mata, C. & Lamattina, L. 2001. Nitric oxide induces stomatal closure and enhances the adaptive plant responses against drought stress. *Plant Physiology* 126(3): 1196-1204.
- Guttieri, M., Bowen, D., Dorsch, J.A., Raboy, V. & Souza, E. 2004. Identification and characterization of a low phytic acid wheat. *Crop Science* 44(2): 418-424.
- Ibrahim, A.M.H. & Quick, J.S. 2001. Genetic control of high temperature tolerance in wheat as measured by membrane thermal stability. *Crop Science* 41(5): 1405-1407.
- Jaleel, C.A., Manivannan, P., Lakshmanan, G.M.A., Gomathinayagam, M. & Panneerselvam, R. 2008. Alterations in morphological parameters and photosynthetic pigment responses of *Catharanthus roseus* under soil water deficits. *Colloids and Surfaces B: Biointerfaces* 61(2): 298-303.
- Javed, I. 2012. Genetics of some potential parameters in *Zea mays* L. under normal and moisture deficit conditions, University of Agriculture, Faisalabad (Unpublished).
- Kashiwagi, J., Krishnamurthy, L., Upadhyaya, H.D., Krishna, H., Chandra, S., Vadez, V. & Serraj, R. 2005. Genetic variability of drought-avoidance root traits in the mini-core germplasm collection of chickpea (*Cicer arietinum* L.). *Euphytica* 146(3): 213-222.
- Khan, N.H., Ahsan, M., Saleem, M. & Ali, A. 2014. Genetic association among various morpho-physiological traits of *Zea mays* under drought condition. *Life Science Journal* 11: 112-122.
- Khodarahmpour, Z. & Hamidi, J. 2011. Evaluation of drought tolerance in different growth stages of maize (*Zea mays* L.) inbred lines using tolerance indices. *African Journal of Biotechnology* 10(62): 13482-13490.
- Kirigwi, F.M., Ginkel, M.V., Trethowan, R., Sears, R.G., Rajaram, S. & Paulsen, G.M. 2004. Evaluation of selection strategies for wheat adaptation across water regimes. *Euphytica* 135(3): 361-371.
- Kitajima, K. & Fenner, M. 2000. Ecology of seedling regeneration. In *Seeds, the Ecology of Regeneration in Plant Communities*, edited by Fenner, M. Oxfordshire: CABI. pp. 331-359.
- Meeks, M., Murray, S.C., Hague, S. & Hays, D. 2013. Measuring maize seedling drought response in search of tolerant germplasm. *Agronomy* 3(1): 135-147.
- Mehdi, S.S., Ahmad, N. & Ahsan, M. 2001. Evaluation of S1 maize (*Zea mays* L.) families at seedling stage under drought conditions. *Online Journal of Biological Sciences* 1: 4-6.
- Mustafa, H.S.B., Ahsan, M., Aslam, M., Ali, Q., Bibi, T., Hasan, E. & Mehmood, T. 2013. Genetic variability and traits association in maize (*Zea mays* L.) accessions under drought stress. *Journal of Agricultural Research* 51(3): 231-238.
- Neelima, S. & Reddy, V.C. 2008. Genetic parameters of yield and fibre quality traits in American cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research* 42(1): 67-70.

- Nour, A.E.M. & Weibel, D.E. 1978. Evaluation of root characteristics in grain sorghum. *Agronomy Journal* 70(2): 217-218.
- Nzuve, F., Githiri, S., Mukunya, D.M. & Gethi, J. 2014. Genetic variability and correlation studies of grain yield and related agronomic traits in maize. *Journal of Agricultural Science* 6(9): 166-176.
- Qayyum, A., Razaq, A., Ahmad, M. & Jenks, M.A. 2011. Water stress causes differential effects on germination indices, total soluble sugar and proline content in wheat (*Triticum aestivum* L.) genotypes. *African Journal of Biotechnology* 10(64): 14038-14045.
- Rezaeieh, K.A. & Eivazi, A. 2011. Evaluation of morphological characteristics in five Persian maize (*Zea mays* L.) under drought stress. *African Journal of Agricultural Research* 6(18): 4409-4411.
- Seghatoleslami, M.J., Kafi, M. & Majidi, E. 2008. Effect of drought stress at different growth stages on yield and water use efficiency of five proso millet (*Panicum miliaceum* L.) genotypes. *Pakistan Journal of Botany* 40(4): 1427-1432.
- Shrimali, M. 2001. Studies on morphological parameters contributing to drought tolerance in cereals. *India New Botanist* 28(1/4): 91.
- Singh, B.D. & Singh, B.D. 1999. *Plant Breeding: Principles and Methods*. New Delhi: Kalyani Publishers.
- Taiz, L. & Zeiger, E. 2006. Stress physiology. In *Plant Physiology*. 4th ed. Massachusetts: Sinauer Associates, Inc.
- Thiry, A.A., Dulanto, P.N.C., Reynolds, M.P. & Davies, W.J. 2016. How can we improve crop genotypes to increase stress resilience and productivity in a future climate? A new crop screening method based on productivity and resistance to abiotic stress. *Journal of Experimental Botany* 67(19): 5593-5603.
- Waqas, M.A., Khan, I., Akhter, M.J., Noor, M.A. & Ashraf, U. 2017. Exogenous application of plant growth regulators (PGRs) induces chilling tolerance in short-duration hybrid maize. *Environmental Science and Pollution Research International* 24(12): 11459-11471. doi: 10.1007/s11356-017-8768-0.
- Wu, Y. & Cosgrove, D.J. 2000. Adaptation of roots to low water potentials by changes in cell wall extensibility and cell wall proteins. *Journal of Experimental Botany* 51(350): 1543-1553.
- Zafar, S.A., Hameed, A., Khan, A.S. & Ashraf, M. 2017. Heat shock induced morpho-physiological response in indica rice (*Oryza sativa* L.) at early seedling stage. *Pakistan Journal of Botany* 49(2): 453-463.
- Zafar, S.A., Shokat, S., Ahmed, H.G.M., Khan, A., Ali, M.Z. & Atif, R.M. 2015. Assessment of salinity tolerance in rice using seedling based morpho-physiological indices. *Advancement in Life Sciences* 2(4): 142-149.
- Fahad Masoud Wattoo & Rashid Mehmood Rana  
Department of Plant Breeding & Genetics  
PMAS-Arid Agriculture University  
Rawalpindi  
Pakistan
- Sajid Fiaz\*  
State Key Laboratory of Rice Biology  
China National Rice Research Institute  
Hangzhou 310006  
China
- Syed Adeel Zafar  
National Key Facility of Crop Gene Resources and Genetic Improvement  
Institute of Crop Science  
Chinese Academy of Agricultural Sciences  
Beijing 100081  
China
- Mehmood Ali Noor & Shoaib ur Rehman  
Institute of Crop Science  
Chinese Academy of Agricultural Sciences  
Key Laboratory of Crop Physiology and Ecology  
Ministry of Agriculture  
Beijing 100081  
China
- Hafiz Mumtaz Hassan  
Nuclear Institute for Agriculture and Biology  
Jhang Rd, Faisalabad  
Pakistan
- Muhammad Husnain Bhatti  
Ayub Agriculture Research Institute  
Jhang Rd, Faisalabad  
Pakistan
- Galal Bakr Anis  
Field Crop Research Institute  
Agriculture Research Centre  
Egypt
- Rai Muhammad Amir  
Institute of Food and Nutritional Sciences  
PMAS-Arid Agriculture University Rawalpindi  
Pakistan

\*Corresponding author; email: fiazsajid05@yahoo.com

Received: 28 May 2017  
Accepted: 14 August 2017