

ESTIMATE ON THE SECOND HANKEL FUNCTIONAL FOR FUNCTIONS WHOSE DERIVATIVE HAS A POSITIVE REAL PART

(Anggaran Fungsian Hankel Kedua bagi Fungsi dengan
Terbitan Bahagian Nyata Positif)

AINI JANTENG, SUZEINI ABDUL HALIM & MASLINA DARUS

ABSTRACT

Denote S to be the class of functions which are analytic, normalised and univalent in the open unit disc $D = \{z : |z| < 1\}$. Also denote R to represent the subclass of S whose derivative has a positive real part. Next, write $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ where a_n is a complex constant and denote the q th Hankel determinant for f as $H_q(n)$ for $q \geq 1, n \geq 1$. Our intention is to seek sharp upper bounds for $H_2(n)$, however, we begin by first looking at $H_2(2)$. In this paper, we give the upper bound for the second Hankel determinant for this particular class of functions.

Keywords: Hankel determinant; upper bound

ABSTRAK

Lambangkan S sebagai kelas fungsi yang terdiri daripada fungsi-fungsi analisis, ternormal dan univalen di dalam cakera unit terbuka $D = \{z : |z| < 1\}$. Lambangkan juga R sebagai subkelas bagi S yang terdiri daripada fungsi-fungsi dengan terbitannya mempunyai bahagian nyata yang positif. Seterusnya dapat diungkapkan bahawa $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ dengan a_n adalah pemalar kompleks dan lambangkan penentu Hankel ke- q bagi f sebagai $H_q(n)$ untuk $q \geq 1, n \geq 1$. Makalah ini bertujuan untuk mendapatkan batas atas terbaik bagi $H_2(n)$, namun, ianya dimulakan dengan melihat $H_2(2)$. Dalam makalah ini, batas atas bagi penentu Hankel kedua diberikan.

Kata kunci: penentu Hankel; batasan atas

1. Introduction

Let S denote the class of normalised analytic univalent functions f of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1)$$

where $z \in D = \{z : |z| < 1\}$. In Noonan and Thomas (1976), the q th Hankel determinant for $q \geq 1$ and $n \geq 1$ is stated by Noonan and Thomas as

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q+1} \\ a_{n+1} & \cdots & & \vdots \\ \vdots & & & \\ a_{n+q-1} & \cdots & & a_{n+2q-2} \end{vmatrix}.$$

This determinant has also been considered by several authors. For example, Noor (1983) determined the rate of growth of $H_q(n)$ as $n \rightarrow \infty$ for functions f given by Eq. (1) with bounded boundary. Ehrenborg (2000) studied the Hankel determinant of exponential polynomials and in (Layman 2001), the Hankel transform of an integer sequence is defined and some of its properties discussed by Layman.

Easily, one can observe that the Fekete and Szegő functional is $H_2(1)$. Fekete and Szegő then further generalised the estimate $|a_3 - \mu a_2^2|$ where μ is real and $f \in S$. For our discussion in this paper, we consider the Hankel determinant in the case of $q = 2$ and $n = 2$,

$$H_2(2) = \begin{vmatrix} a_2 & a_3 \\ a_3 & a_4 \end{vmatrix}.$$

We seek upper bound for the functional $|a_2 a_4 - \mu a_3^2|$ where μ is real for functions f belongs to the class R . The class R is defined as follows:

Definition 1.1 Let f be given by Eq. (1). Then $f \in R$ if and only if

$$\operatorname{Re} \{f'(z)\} > 0, \quad (z \in D). \tag{2}$$

We first state some preliminary lemmas, required for proving our result.

2. Preliminary results

Let P be the family of all functions p analytic in D for which $\operatorname{Re} p(z) > 0$ and

$$p(z) = 1 + c_1 z + c_2 z^2 + \dots \tag{3}$$

for $z \in D$.

Lemma 2.1 (Pommerenke (1975)) *If $p \in P$ then $|c_k| \leq 2$ for each k .*

Lemma 2.2 (Grenander & Szegő (1958)) *The power series for p given in Eq. (3) converges in D to a function in P if and only if the Toeplitz determinants*

$$D_n = \begin{matrix} \left| \begin{array}{cccc} 2 & c_1 & c_2 & \dots & c_n \\ c_{-1} & 2 & c_1 & \dots & c_{n-1} \\ M & & & & \\ c_{-n} & c_{-n+1} & c_{-n+2} & \dots & 2 \end{array} \right|, & n = 1, 2, 3, \dots \end{matrix} \quad (4)$$

and $c_{-k} = \bar{c}_k$, are all non negative. They are strictly positive except for $p(z) = \sum_{k=1}^m \rho_k p_0(e^{it_k z})$, $\rho_k > 0$, t_k real and $t_k \neq t_j$ for $k \neq j$; in this case $D_n > 0$ for $n < m - 1$ and $D_n = 0$ for $n \geq m$.

This necessary and sufficient condition is due to Carathéodory and Toeplitz and can be found in (Grenander & Szegö (1958)).

3. Main results

Theorem 3.1 Let $f \in R$. Then

$$|a_2 a_4 - \mu a_3^2| \leq \begin{cases} \frac{(27-16\mu)^2}{144(9-8\mu)} - \frac{4\mu}{9}, & \text{if } \mu \leq 0, \\ \frac{(27-32\mu)^2}{144(9-8\mu)} + \frac{4\mu}{9}, & \text{if } 0 \leq \mu \leq \frac{27}{32}, \\ \frac{4\mu}{9}, & \text{if } \frac{27}{32} \leq \mu \leq \frac{27}{16}, \\ \frac{(16\mu-27)^2}{144(8\mu-9)} + \frac{4\mu}{9}, & \text{if } \mu \geq \frac{27}{16}. \end{cases}$$

Proof: Since $f \in R$, it follows from Eq. (2) that

$$f'(z) = p(z) \quad (5)$$

for some $z \in D$. Equating coefficients in Eq. (5), yields

$$\left. \begin{array}{l} 2a_2 = c_1 \\ 3a_3 = c_2 \\ 4a_4 = c_3 \end{array} \right\}. \quad (6)$$

From Eq. (6),

$$|a_2 a_4 - \mu a_3^2| = \left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right|. \quad (7)$$

Lemma 2.2 can then be used to obtain the proper bound on Eq. (7). We may assume without restriction that $c_1 \geq 0$. Rewriting Eq. (4) for the cases $n = 2$ and $n = 3$, result in

$$D_2 = \begin{vmatrix} 2 & c_1 & c_2 \\ c_1 & 2 & c_1 \\ \bar{c}_2 & c_1 & 2 \end{vmatrix} = 8 + 2 \operatorname{Re}\{c_1^2 c_2\} - 2|c_2|^2 - 4c_1^2 \geq 0,$$

which is equivalent to

$$2c_2 = c_1^2 + x(4 - c_1^2) \quad (8)$$

for some $x, |x| \leq 1$.

Further, $D_3 \geq 0$ is equivalent to

$$\left| (4c_3 - 4c_1 c_2 + c_1^3)(4 - c_1^2) + c_1(2c_2 - c_1^2)^2 \right| \leq 2(4 - c_1^2)^2 - 2|2c_2 - c_1^2|^2;$$

and this, with Eq. (8), provides the relation

$$4c_3 = c_1^3 + 2(4 - c_1^2)c_1 x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z, \quad (9)$$

for some value of $z, |z| \leq 1$.

Suppose now that $c_1 = c$ and $0 \leq c \leq 2$. Using Eq. (8) along with Eq. (9), we obtain

$$\begin{aligned} & \left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right| \\ &= \left| \frac{(9-8\mu)c^4}{288} + \frac{(9-8\mu)(4-c^2)c^2 x}{144} + \frac{(4-c^2)(1-|x|^2)cz}{16} - \frac{(4-c^2)x^2(c^2(9-8\mu)+3-\mu)}{288} \right| \end{aligned}$$

Triangle inequality gives

$$\begin{aligned} & \left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right| \\ & \leq \frac{|9-8\mu|c^4}{288} + \frac{c(4-c^2)}{16} + \frac{|9-8\mu|c^2(4-c^2)\rho}{144} + \frac{(4-c^2)(|9-8\mu|c^2+32|\mu|-18c)\rho^2}{288} \\ & = F(\rho) \end{aligned} \quad (10)$$

with $\rho = |x| \leq 1$. This gives rise to

$$F'(\rho) = \begin{cases} \frac{(9-8\mu)(4-c^2)c^2}{144} + \frac{(4-c^2)\{(9-8\mu)c^2-32\mu-18c\}\rho}{144}, & \text{if } \mu \leq 0, \\ \frac{(9-8\mu)(4-c^2)c^2}{144} + \frac{(4-c^2)\{(9-8\mu)c^2+32\mu-18c\}\rho}{144}, & \text{if } 0 \leq \mu \leq \frac{9}{8}, \\ \frac{(8\mu-9)(4-c^2)c^2}{144} + \frac{(4-c^2)\{(8\mu-9)c^2+32\mu-18c\}\rho}{144}, & \text{if } \mu \geq \frac{9}{8}. \end{cases}$$

and again for all the cases above, $F'(\rho) > 0$ for $\rho > 0$; implying that F is an increasing function with $\text{Max}_{\rho \leq 1} F(\rho) = F(1)$.

Now let

$$G(c) = F(1) = \frac{|9-8\mu|c^4}{288} + \frac{c(4-c^2)}{16} + \frac{|9-8\mu|c^2(4-c^2)}{144} + \frac{(4-c^2)\{|9-8\mu|c^2+32|\mu|-18c\}}{288}. \quad (11)$$

(i) First, let us consider the case $\mu \leq 0$.

Eq. (11) gives

$$G'(c) = \frac{c}{36} \{ -(9-8\mu)c^2 + 27 - 16\mu \}.$$

Elementary calculation reveals that G attains its maximum value at $c = \sqrt{\frac{27-16\mu}{9-8\mu}}$.

The upper bound for Eq. (10) corresponds to $\rho = 1$ and $c = \sqrt{\frac{27-16\mu}{9-8\mu}}$, in which case

$$\left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right| \leq \frac{(27-16\mu)^2}{144(9-8\mu)} - \frac{4\mu}{9}.$$

(ii) Secondly, we consider the case $0 \leq \mu \leq \frac{27}{32}$.

This gives

$$G'(c) = \frac{c}{36} \left\{ -(9-8\mu)c^2 + 27 - 32\mu \right\}$$

where G attains its maximum value at $c = \sqrt{\frac{27-32\mu}{9-8\mu}}$. Hence, we obtain

$$\left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right| \leq \frac{(27-32\mu)^2}{144(9-8\mu)} + \frac{4\mu}{9}.$$

(iii) To prove the third result, we consider two cases.

First, assume that $\frac{27}{32} \leq \mu \leq \frac{9}{8}$.

In this case, G attains its maximum value at $c = \sqrt{\frac{32\mu-27}{9-8\mu}}$.

Next, consider the case $\frac{9}{8} \leq \mu \leq \frac{27}{16}$. For this case, Eq. (11) gives rise to

$$G'(c) = \frac{c}{36} \left\{ -(8\mu-9)c^2 + 16\mu - 27 \right\},$$

where G attains its maximum value at $c = \sqrt{\frac{27-16\mu}{8\mu-9}}$.

In both cases, the upper bound is attained as

$$\left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right| \leq \frac{4\mu}{9}.$$

(iv) Finally, consider $\mu \geq \frac{27}{16}$.

Here, G attains its maximum value at $c = \sqrt{\frac{16\mu-27}{8\mu-9}}$. Hence,

$$\left| \frac{c_1 c_3}{8} - \frac{\mu c_2^2}{9} \right| \leq \frac{(16\mu - 27)^2}{144(8\mu - 9)} + \frac{4\mu}{9}.$$

This completes the proof of theorem. \square

References

- Ehrenborg R. 2000. The Hankel determinant of exponential polynomials. *American Mathematical Monthly* **107**: 557-560.
- Fekete M. & Szegő G. 1933. Eine Bemerkung über ungerade schlichte Funktionen. *J. London Math. Soc.* **8**: 85-89.
- Grenander U. & Szegő G. 1958. *Toeplitz forms and their application*. Berkeley: Univ. of California Press.
- Layman J.W. 2001. The Hankel transform and some of its properties. *J. of Integer Sequences* **4**: 1-11.
- Noonan J.W. & Thomas D.K. 1976. On the second Hankel determinant of areally mean p-valent functions. *Trans. Amer. Math. Soc.* **223**(2): 337-346.
- Noor K.I. 1983. Hankel determinant problem for the class of functions with bounded boundary rotation. *Rev. Roum. Math. Pures Et Appl.* **28**(8): 731-739.
- Pommerenke Ch. 1975. *Univalent functions*. Göttingen: Vandenhoeck and Ruprecht.

School of Science and Technology
Universiti Malaysia Sabah
Locked Bag No. 2073
88999 Kota Kinabalu, Sabah
MALAYSIA
*E-mail : aini_jg@ums.edu.my**

Institute of Mathematical Sciences
Faculty of Science
Universiti Malaya
50603 Kuala Lumpur
MALAYSIA
E-mail: suzeini@um.edu.my

School of Mathematical Sciences
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
43600 Bangi, Selangor D.E.
MALAYSIA
E-mail: maslina@ukm.my

* Corresponding author