

TRANSPOSABLE ELEMENTS AND THE DETECTION OF SOMACLONAL VARIATION IN PLANT TISSUE CULTURE: A REVIEW

AZMAN, A.S.¹, MHIRI, C.², GRANDBASTIEN, M.A.² and TAM, S.M.^{3*}

¹Jeffrey Cheah School of Medicine and Health Sciences,
Monash University Malaysia, Jalan Lagoon Selatan, 46150, Bandar Sunway, Selangor, Malaysia

²Institut Jean-Pierre Bourgin, UMR1318 INRA-AgroParis Tech,
Batiment 2, INRA Versailles, Route de St Cyr, 78026 Versailles cedex, France

³School of Biosciences, Taylor's University, Jalan Taylor's,
47500, Subang Jaya, Selangor, Malaysia

*E-mail: shehmay.tam@taylors.edu.my

ABSTRACT

Plant tissue culture uses the genetic potential, or totipotency, of plants to regenerate and give rise to a whole plant. Tissue culture-derived plants are expected to have identical genetic material to the parent, and thus can serve as an effective tool for controlled, mass clonal propagation. However, somaclonal variation has been observed in plant tissue culture due to either point mutations, transposition activity of mobile genetic elements, chromosomal rearrangements, or ploidy level changes, causing genetic instability. Tissue culture-induced mutations associated with transposable element activities have been reported from many plant studies, related to stress conditions during tissue culture such as wounding, exposure to hormones and/or specific compounds in the growth media, and genomic shocks from cytological changes. Transposable elements are repetitive DNA fragments with the ability to transpose from one region to another within a genome. Mutations that may occur during plant tissue culture suggest that the plantlets have to be regularly monitored. Methyl-sensitive transposon display (MSTD) is an advanced, efficient DNA fingerprinting technique that can simultaneously detect genetic variation, changes in transposable element insertion sites and the status of cytosine methylation of DNA in plant genomes. The main principles and notes for application of MSTD such as design of element-specific primers, identification of transposable element sequences in plant genomes, selection of the isochizomer enzymes used, and different classes of banding pattern shown are briefly discussed based on our preliminary work with *Nicotiana benthamiana* (Tnt1 retroelement) and *Musa acuminata* (Copia-33 Mad-I retroelement) examples.

Key words: Methyl-sensitive transposon display (MSTD), somaclonal variation, transposable elements

INTRODUCTION

Plant tissue culture, also known as plant cell, sterile, axenic or *in vitro* culture, is a technique of growing plant cells, tissues or organ in an artificial gel or liquid media supplemented with nutrients, vitamins and plant growth regulators under controlled and sterile conditions (Singh and Kumar, 2009). Plant cells possess sufficient genetic potential to be able to regenerate and give rise to a whole plant (totipotency) – making plant tissue culture an important method in plant biotechnology studies as well as having commercial applications (Thorpe, 2007). Originally proposed by Gottlieb Haberlandt in 1902, single cells were predicted to regenerate

into a complete and functional plant (Krikorian and Berquam, 1969), this hypothesis of totipotency was first proved using tobacco cells (Vasil and Hildebrand, 1965) and later using tobacco protoplasts to regenerate a new plant (Takebe *et al.*, 1971).

Plants grown via tissue culture are expected to have identical genetic material to the parent and thus, keep their intrinsic characteristics. This method of establishing genetically identical clones from an organism's tissue, capable of generating into a complete plant is known as clonal propagation or micropropagation (Vizel *et al.*, 2010). Micropropagation has been extensively used for the multiplication and growth of many plants in horticulture and agriculture, ranging from wild species such as orchids, ferns, carnivorous plants,

* To whom correspondence should be addressed.

medicinal plants, woody plants, and cultivated crops (Fay, 1992; Debnath *et al.*, 2006; Gatica-Arias *et al.*, 2008; Dam *et al.*, 2010; Vergara-Galicia *et al.*, 2010; Paek *et al.*, 2011). Main advantages associated with micropropagation include the large number of homogenous and genetically stable plants that can be produced under sterile condition, relative ease of controlling the environmental and nutritional conditions and rapid production of new varieties (Dobráński and Texeira de Silva, 2010). Two commonly used methods in micropropagation include organogenesis and somatic embryogenesis (Rout *et al.*, 2006). These techniques share similar main aims of enhancing the rate of multiplication of regenerated plants and the production of disease free plants (Eka *et al.*, 2005).

Mutation leads to phenotypic, genetic variation and genome instability

Plant tissue culture can be seen as an effective tool for the large scale, controlled production of plant material and represent an option for mass clonal propagation (Chandrika and Rai, 2009). Even though plant tissue culture provides many advantages, one major drawback associated with it is mutations that can cause phenotypic and genetic changes i.e. somaclonal variation. Mutations can cause phenotypic alteration, genetic variation and genome instability among the regenerated plants from a single donor clone. Possible mechanisms that may cause a mutation to occur during the tissue culture process are under the controls of genetic and epigenetic systems where it can affect the genetic and genome stability of the plant (Temel *et al.*, 2008). Somaclonal variation may arise due to point mutations, the activation of mobile genetic elements, chromosomal rearrangements, or ploidy level changes (Jaligot *et al.*, 2000). Any genetic changes induced by tissue culture condition will probably produce a plant with unique heritable characteristics (Soniya *et al.*, 2001). Even though the resulting mutant phenotypes of the plants (e.g. alteration in leaf shape, dwarfing, and other changes in growth habit) are unique compared to the mother plant, they are not normally deemed useful in micropropagation or for crop improvement (Evans, 1989).

Mutations are induced by specific components of culture medium where the unnatural condition and environment, and hormone supplemented in the media can generate spontaneous and heritable genetic changes due to a “shocking” experience of isolation. Rates of genetic instability (mutations) are dependent on the type and quantity of the plant growth regulators, such as auxin, cytokinin and abscisic acid in the media (Michiba *et al.*, 2001), the duration of the regenerated plant in the culture (Fras and Maluszynska, 2004) and the degree of

endopolyploidy of the explants (Nontaswatsri and Fukai, 2005). Typically, high number of subcultures, exposure to high concentration of plant growth regulators and long term culture are factors that can result in mutations (Marum *et al.*, 2009). Epigenetic changes are not due to sequence modification and chromosomal aberration (Akimoto *et al.*, 2007). Tsiftaris *et al.* (2005) defined epigenetic changes as “all meiotically and mitotically heritable changes in gene expression that are not coded in the DNA sequence itself”. This aspect commonly occurred when the plants are under stress conditions – in tissue culture environments such as prolonged culture time and exposure to plant growth regulators (Vázquez and Linacero, 2010). One interesting source of genomic instability is linked to the activation of transposable elements that might be responsible for tissue-culture induced mutation (Momose *et al.*, 2010; Karajol and Naik, 2011; Bui and Grandbastien, 2012).

Activation of transposable elements during plant tissue culture

Mobile genetic elements or transposable elements (TEs) are repetitive DNA fragments that have the ability to move or transpose from one area to another area within a genome (Lisch, 2009). Transposable elements contribute to the size of plant genomes; for example, in certain species of grass, they may comprise of 50% up to 80% of the genomes (Meyers *et al.*, 2001; Piegu *et al.*, 2006; Dooner and Weil, 2007; Macas *et al.*, 2007). Shirasu *et al.* (2000) reported that about 40% to 70% of the total DNA content of crop plants may be made up of repetitive elements, and it was reported that a correlation exists between the evolution of LTR retrotransposons and genome size (Zedek *et al.*, 2010). Transposable elements may greatly impact host genomes by causing mutations when they transpose into or near a gene (Chen *et al.*, 2005). They can alter expression of a nearby gene by operating as promoters, silencers, enhancers, or act as targets of epigenetic modification and other alternative splicing events (Kazazian, 2004; Bui and Grandbastien, 2012). They can also serve as locations for homologous recombination, resulting in chromosomal rearrangements such as deletions, inversions, duplications or translocations due to high, repetitive copies of integrated elements (populations of the same family of elements) existing in the genome (Devos *et al.*, 2002; Kolomietz *et al.*, 2002; Bailey *et al.*, 2003; Sen *et al.*, 2006).

Generally divided into two classes based mainly on their transposition intermediates, retrotransposons (also known as Class I elements) are elements which move via an RNA intermediate while DNA transposons (or Class II elements) directly transpose in the form of DNA (Feschotte *et*

al., 2002; Todorovska, 2007). Retrotransposons represent a substantial percentage of eukaryotic genomes as they transpose by a “copy and paste” mechanism catalyzed by reverse transcriptase and endonuclease (EN) domains of a polyprotein encoded by themselves or by other retrotransposons (Collier and Largaespada, 2007). Based on their structure, retrotransposons can be further divided into two major types – elements that are flanked by long terminal repeats (LTRs) and non-LTR retrotransposon. Non-LTR retrotransposons are divided into long interspersed nuclear elements (LINEs) and short interspersed nuclear elements (SINEs) (Feschotte *et al.*, 2002; Karajol and Naik, 2011). LINEs are autonomous retrotransposons that encode the enzymatic machinery required for their propagation (Ostertag and Kazazian, 2001) while SINEs are non-autonomous and require the enzymatic machinery of LINE elements for retrotransposition (Lenoir *et al.*, 2001; Ostertag *et al.*, 2003).

In general, DNA transposons move only as DNA from one genomic area to another and can be deleted precisely though at a relatively low frequency (Huang *et al.*, 2008). Mobility of these elements is performed by a “cut-and-paste” mechanism using an element encoded transposase (TPase) (Arkhipova and Meselson, 2005). However, more recent work interestingly revealed that DNA transposons may be further divided into three major classes which consist of (i) “cut-and-paste” DNA transposons, (ii) rolling-circle DNA transposons or Helitrons, which were found to use a replicative transposition mode and (iii) self-synthesizing DNA transposon or Polintons. DNA transposons are characterized by a superfamily-specific transposase core which differs between superfamilies (Kapitonov and Jurka, 2008). DNA transposons are also distinguished by terminal inverted repeats (TIRs) at their extremities and similar to retroelements, both classes of elements form target site duplication (TSDs) during insertion (Wicker *et al.*, 2007). Upon transposition, the TIRs are identified by the DNA-binding domains of TPases and this element usually excises from one location and reinserts into elsewhere in the genome (Chandler and Mahillon, 2002). Both class I and II transposable elements have been found to contain autonomous and non-autonomous elements, whereby autonomous elements have open reading frames (ORFs) that encode the products required for the whole process of transposition. Non-autonomous elements require the presence of an active autonomous element in order to move since they have no significant coding capacity (Lisch, 2009).

During the process of plant tissue culture, transposable elements have been found to be a significant source of mutations and their activities

can be influenced (silenced) by some epigenetic mechanisms due to changes in the environment and culture condition (Gaut and Ross-Ibarra, 2008). Tissue culture stress is related to wounding during the isolation of explants, exposure to hormones and/or specific compounds in the growth media during *in vitro* cell culture or tissue culture. All these factors may activate transposable elements and result in changes in their copy numbers (Melayah *et al.*, 2001; Kubis *et al.*, 2003; Bui and Grandbastien, 2012). Due to the (potential) highly mutagenic impact of TE activities, it is known that host plants have evolved different epigenetic mechanisms to regulate element activity. These mechanisms include transcriptional silencing via DNA methylation, histone modification and alterations in chromatin packing and condensation, as well as post-transcriptional silencing of elements by the RNAi pathway, and the more recently described transposon - derived small interfering (si) RNAs and RNA directed DNA methylation (RDRM) and histone modification to suppress transpositional activity (Martienssen and Colot, 2001; Lippman and Martienssen, 2004; Matzke and Birchler, 2005; Slotkin and Martienssen, 2007; Ito 2012, 2013).

In plants, DNA (CpG) methylation have been reported to normally repress transpositional activity of diverse transposons, and studies comparing transcriptional and transpositional activities of elements (Evade, ONSEN), using methylation and siRNAs mutants also demonstrated regulation of transposable elements via RNAi-mediated chromatin modifications (Mirouze *et al.*, 2009; Ito *et al.*, 2011). Therefore, normal epigenetic program and re-patterning that may not be established properly in regenerated plants could lead to activation of transposable elements due to hypomethylation in tissue culture conditions (Kaeppeler *et al.*, 2000). However, it is interesting to note that exceptions have been detected, whereby CpG methylation instead elevated transpositional activity of certain elements (Sleeping Beauty, Frog Prince and Minos) of the Tc1/mariner superfamily (Yusa *et al.*, 2004; Ikeda *et al.*, 2007; Jursch *et al.*, 2013). It was also proposed that chromosome breakages that sometimes occurred in tissue culture induce genomic shock which in turn activates the transposable elements, though the main cause of chromosome breakage in plant tissue culture is still unclear but may be related to changes in DNA methylation pattern resulting in chromosome bridges and breakage (Kaeppeler *et al.*, 2000).

Tissue culture-induced mutations associated with transposable element activities have been reported in many plant species - various studies have reported that *in vitro* tissue culture can activate transposable elements such Ac in maize (Brettell and Dennis, 1991), Tto1 in *Nicotiana tabacum*

(Hirochika, 1993), Tos17 (Miyao *et al.*, 2003) and Lullaby (Picault *et al.*, 2009) in *Oryza sativa*, Tnt1 in *Medicago truncatula* (d'Erfuth *et al.*, 2003), and Rtsp-1 in *Ipomea batatas* (Tahara *et al.*, 2004). Hirochika *et al.* (1996) originally showed that Tos retrotransposons were inactive in normal plants but increased in transcription during tissue culture. Kikuchi *et al.* (2003) observed the activation of MITEs (*mPing*) during anther culture of rice, which led to transpositions into new gene locations. Interestingly, Huang *et al.* (2009) reported transposition of *nDaiZ9* was activated in most of the samples upon treatment with a DNA methylation inhibitor (5-azaC). Another study suggested that prolonged tissue culture is needed for more transposition of *Tos17* in rice tissue culture leading to more and severe gene mutations (Kuan *et al.*, 2010). Kour *et al.* (2009) showed that transposable elements had been activated in callus regenerated plants bearing the B chromosome of *Plantago lagopus*, and deduced that activation of transposable elements was highly related to the condition in tissue culture since there were no changes in the B chromosome of the mother plant. Recently, microarrays and next generation sequencing (NGS) analysis detected genome-wide activities of transposable elements in tissue culture and suspension cells of *Arabidopsis thaliana* (Tanurdzic *et al.*, 2008) and tissue culture derived plants of *Oryza sativa* (Picault *et al.*, 2009; Sabot *et al.*, 2011; Miyao *et al.*, 2012).

An advanced molecular marker technique to detect genomic instability: Methyl-Sensitive Transposon Display (MSTD)

Mutations that may occur during plant tissue culture suggest that the plantlets have to be regularly monitored. Various molecular marker techniques are available for use to detect somaclonal variation. Most of the molecular marker techniques involve amplifications of genomic DNA with short random or specific primers. For example, tissue-culture induced mutations of different plant tissue cultures have been screened using Random Amplified Polymorphic DNAs (RAPD) (Eshraghi *et al.*, 2005; Rasheed *et al.*, 2005; Qin *et al.*, 2007; Venkatachalam and Sreedhar 2007; Sianipar *et al.*, 2008; Elmeer *et al.*, 2009), Amplified Fragment Length Polymorphism (AFLP) (Pontaroli and Camadro, 2005; Puente *et al.*, 2008; Chuang *et al.*, 2009), microsatellite markers or simple sequence repeats (SSR) (Morgante *et al.*, 2002; Lopes *et al.*, 2006; Burg *et al.*, 2007), Sequence-Specific Amplified Polymorphism (SSAP) (Venturi *et al.*, 2006; Du *et al.*, 2009; Wegscheider *et al.*, 2009) and Methyl-Sensitive Amplified Polymorphism (MSAP) (Peraza-Echeverria *et al.*, 2001; Jaligot *et al.*, 2004;

Lu *et al.*, 2008). Recently, another modified advanced method called Methyl-Sensitive Transposon Display or MSTD (Figure 1a), was shown to be useful for detecting genetic variation, changes in transposable element banding pattern and DNA methylation all in one technique (Parisod *et al.*, 2009).

Methyl-sensitive transposon display (MSTD) is actually a combination method of MSAP (Reyna-Lopez *et al.*, 1997) and Sequence Specific Amplification Polymorphism (SSAP), which is highly similar to Transposon Display (Casa *et al.*, 2000; Syed and Flavell, 2007). Both MSAP and MSTD are efficient DNA fingerprinting methods to detect the conditions of large scale cytosine methylation (status) in plant genomes, and differences in the banding patterns rapidly reveal and identify changes in methylation between the genomes of the sampled plants. In addition, MSTD is also able to provide information on epigenetic (methylation status) of adjacent regions and population dynamics (copy numbers) of transposable elements (TEs) insertions (Zerjal *et al.*, 2009). The MSTD technique relies on primer(s) anchored at TE extremities to specifically amplify digested genomic DNA of the insertion sites of TEs and therefore, this technique can be used to comparatively evaluate the copy number of TEs. An amplified fragment will consist of sequences of part of the element, and the genomic sequence adjacent to the border of the element - and this varies in length according to the location of the nearest restriction site on the genomic DNA (de Setta *et al.*, 2007; Parisod *et al.*, 2014). Results from MSTD are also useful to potentially provide information on the impact of TE insertions on the genome since certain elements such as retrotransposons are widely distributed throughout the genome (Miyao *et al.*, 2003) and can give insights to improve our understanding, predict and possibly track genomic evolution due to transposition events (Mandal *et al.*, 2006). Parisod *et al.* (2009) performed a study in *Spartina* using three elements, namely *Ins2*, *Cassandra* and *Wis-like*, which were mostly shared by both parents and hybrid plants. Their results found that CpG methylation and major structural changes have occurred near the TE insertion sites and thus caused alterations in the plant genome following inter-specific hybridization and/or genome duplication.

At present, a limited number of studies have been performed to detect genomic instability that may have occurred during plant tissue culture of Malaysian crops. Perhaps one main hindrance is the design of element-specific primers, which requires existing DNA sequence information of well-characterized elements. Nowadays, however, many genomics studies using the application of high-

throughput sequencing technologies have been performed resulting in abundant information on genomic sequences for many crop genomes (for example, see Plant Genomes Central at National Center for Biotechnology Information (NCBI) for current list of plant genomic projects, @ <http://www.ncbi.nlm.nih.gov/genomes/PLANTS/PlantList.html>). Available full genome sequence information (datasets) have even been exploited to develop a method that can predict the transposon insertion display banding pattern of a given transposable element primer (Le and Bureau, 2004) so designed primers can be tested virtually before being applied in MSTD. In addition, there is an existing site hosting the Plant Repeat Databases (<http://plantrepeats.plantbiology.msu.edu/about.html>), created to assist in the compilation and identification of repeat sequences in plant genomes whereby repetitive DNA sequences of selected plant genera are regularly queried and compiled from GenBank and other published records (Ouyang and Buell, 2004).

Our own preliminary work testing MSTD on the model plant *Nicotiana benthamiana* and a crop plant, banana (*Musa acuminata*) showed that it was relatively easy to adopt the MSTD protocol. The primer used for *Nicotiana* was based on the well-studied tobacco retroelement Tnt1, which is one of the few active retrotransposons in plants (Melayah *et al.*, 2001). Transposition of this element has been shown previously to be activated during protoplast culture in tobacco and tissue culture in the heterologous hosts *Arabidopsis thaliana* and *Medicago truncatula* during the early steps of the *in vitro* transformation-regeneration process (Grandbastien *et al.*, 1998; d'Erfurth *et al.*, 2003). Sequences used to generate a retroelement primer for *Musa acuminata* were directly obtained from NCBI whereby a quick search revealed the availability of a large Bacterial Artificial Chromosome (BAC) clone sequence (AC226035). The program Repeat Masker, Genetic Information Research Institute (GIRI) (Kohany *et al.*, 2006) identified several transposable elements of which one was named Copia-33 Mad-I, similar to *copia* elements from *Malus domestica*. The primer used was then designed using Primer3 Input version 0.4.0 (Rozen and Skaletsky, 2000), positioned facing outwards on the long terminal repeat (5' LTR) sequence of the Copia-33 Mad-I retrotransposon element.

Our test application of MSTD on *Nicotiana* (normal and mutant phenotype "grassy") and banana tissue cultured plantlets (normal and mutant phenotype "cabbage-like") with the Tnt1 and Copia-33 Mad-I primers respectively produced results as shown in Figure 1(b) and 1(c). Results demonstrated banding pattern differences between

the normal and mutant plantlets in both the *Nicotiana* and banana samples (Azman, 2011). Polymorphic bands can be seen between normal and mutant plantlets, as well as between the samples digested by *MspI* and *HpaII*, by the disappearance and appearance of bands, indicating differences in DNA methylation pattern occurring near the retroelement insertion sites or element activity, in both normal as well as mutated plants. In addition, the MSTD gel results also demonstrated that the genome of banana contained many copies of Copia-33 Mad-I, suggesting that this retrotransposon might be present in high copy number and thus possibly play a role in the evolution of the banana genome.

A notable step in the MSTD technique which could influence the success of the analysis is selection of the isochizomer enzymes used for detection of methylation status. Isochizomers *MspI* and *HpaII* have been widely used for methyl-sensitive displays since these are frequent cutters compared to other isochizomer enzymes such as *MseI* and *TaqI* (Takata *et al.*, 2007). Both *MspI* and *HpaII* recognize the same tetranucleotide sequence of 5'-CCGG-3' but have different sensitivity to methylation (Krauss *et al.*, 2009). *HpaII* activity is blocked when either internal or external cytosine is fully methylated (both strands) while *MspI* is only blocked when the external cytosine is fully methylated or in a rarer case, when the sequence is hemi-methylated (Xiong *et al.*, 1999). Different levels of methylation of the internal cytosine will result in different cutting by the restriction enzymes thus generating uneven PCR bands between two digests (Zerjal *et al.*, 2009). Hence, the quality of extracted genomic DNA (gDNA) used for the digestion process is an important aspect to ensure the success of the protocol. Significant contamination occurring in the gDNA samples may affect the ability of the restriction enzyme to digest the gDNA as the presence of proteins, polysaccharides and/or polyphenols may interfere with restriction enzyme activity during the digestion process, as well as *Taq polymerase* activity during PCR amplification (Angeles *et al.*, 2005).

The MSTD analysis can provide a powerful technique to investigate genomic changes around transposable elements that occurred during the plant tissue culture process. Three classes of bands may be identified from the polyacrylamide gel electrophoresis (PAGE) results – the first class represent un-methylated bands, present in both *MspI*- and *HpaII*- digested gDNAs; a second class with presence of bands in gDNA digested with *MspI* but missing in *HpaII* digested gDNA, representing methylated sequences and a rare third class of bands present in *HpaII* digested gDNA but missing from




Fig. 1. (a) Schematic representation depicting the principle of methyl-sensitive transposon display (MSTD) adapted from Parisod *et al.* (2014). Digestion of genomic DNA with methylation sensitive isoschizomers (such as *MspI* and *HpaII*) is followed by digestion with a frequent cutter such as *EcoRI*, and then adaptors are ligated to the DNA fragments. PCR amplification using a specific transposable element anchored primer with the complementary adaptor primer will generate bands that are separated by polyacrylamide gel electrophoresis. (b) An example of MSTD profiles generated by using the restriction enzymes *MspI* (C2 and C3) and *HpaII* (C2' and C3') of tissue culture-derived samples of normal (C3) and mutant "grassy" (C2) *Nicotiana benthamiana*. (c) Preliminary example of MSTD profiles generated by using the restriction enzymes *MspI* (2.1, 2.3 and 3.3) and *HpaII* (2.1', 2.3' and 3.3') of tissue culture-derived samples of normal (2.1) and mutant "cabbage" (2.3 and 3.3) *Musa acuminata*.

the *MspI* digested gDNA – these bands might indicate hemi-methylation of the external cytosine (Zerjal *et al.*, 2009; Parisod *et al.*, 2014). Presence of a specific band from a selected sample may correspond to either a transposition event or demethylation of the area near a TE insertion. However, CpCpG methylation on both DNA strands prevents the enzymes from digesting the DNA, therefore, this technique does not perform well in regions of the genome that are heavily methylated. An absence of a band may signify a change in TE insertion or increased levels of methylation and this should be recognized during interpretation of the MSTD profiles (Parisod *et al.*, 2014). Protocol adjustments and further explanations to aid the interpretation of the banding patterns produced by the MSAP method, which is similar to MSTD, is available for samples that display difficult, complex patterns (Fulneček and Kovařík, 2014).

CONCLUSION

The occurrence of mutations during plant tissue culture can be a problem as it leads to plant genomic instability. As a result, the plantlet will possess somaclonal variation and may display different phenotypes from the mother plant. Studies which can offer insights into the causes of mutations are important in order to better understand the types of mutation that may happen during the *in vitro* process so that improvement on the procedure can be done to reduce the number of mutants. Transposable elements constitute a large fraction of repetitive regions in plant genomes and its mobilization affect the evolution of current genomes. Tissue culture conditions have been observed in many plant studies to induce transposable element activities, making elements a source of mutations during tissue culture. An advanced molecular marker technique such as MSTD has not only been proven efficient for the detection of genomic changes in mutant plants and able to detect the complexity of DNA methylation changes during plant development, it also offers an opportunity to study the effect(s) of transposition of TEs by examining insertions potentially associated with specific phenotypic changes of the plant. As a start, this can be done as unique or specific MSTD bands can be isolated, cloned and sequenced in order to discover the affected DNA regions in the host plant. Further genetic studies can then be performed to follow the inheritance (segregation) of the specific marker with the phenotype to confirm the genotype-phenotype association.

ACKNOWLEDGEMENTS

We would like to thank Dr. Norzihan Abdullah from the Malaysian Agricultural Research and Development Institute (MARDI) for providing the banana tissue culture plantlets for the preliminary study.

REFERENCES

- Akimoto, K., Katakami, H., Kim, H.J., Ogawa, E., Sano, C.M., Wada, Y. and Sano, H. (2007). Epigenetic inheritance in rice plants. *Annals of Botany* **100**: 205-217.
- Angeles, J.G.C., Laurena, A.C. and Tecson-Mendoza, E.M. (2005). Extraction of genomic DNA from lipid-, polysaccharide-, and polyphenol-rich coconut (*Cocos nucifera* L.). *Plant Molecular Biology Reporter* **23**: 297a-297i.
- Arhipova, I.R. and Meselson, M. (2005). Diverse DNA transposons in rotifers of the class *Bdelloidea*. *Proceeding of National Academy of Science of United States of America* **102(33)**: 11781-11786.
- Azman, A.S. (2011). Detection of genomic instability during somatic embryogenesis of oil palm. Honours thesis, Monash University Sunway Campus.
- Bailey, J.A., Liu, G. and Eichler, E.E. (2003). An Alu transposition model for the origin and expansion of human segmental duplication. *The American Journal of Human Genetics* **73**: 823-834.
- Brettell, R.I.S. and Dennis, E.S. (1991). Reactivation of a silent Ac following tissue culture is associated with heritable alterations in its methylation pattern. *Molecular Genomics and Genetics* **229**: 365-372.
- Bui, Q.T. and Grandbastien, M.-A. (2012). LTR Retrotransposons as controlling elements of genome response to stress? In: M.-A. Grandbastien and J.M. Casacuberta (eds.). *Plant Transposable Elements. Topics in Current Genetics* 24, Springer-Verlag, Berlin Heidelberg. pp. 273-296.
- Burg, K., Helmersson, A., Bozhkov, P. and von Arnold, S. (2007). Development and genetic variation in nuclear microsatellite stability during somatic embryogenesis in pine. *Journal of Experimental Botany* **58(3)**: 687-698.
- Casa, A.M., Brouwer, C., Nagel, A., Wang, L., Zhang, Q., Kresovich, S. and Wessler, S.R. (2000). The MITE family heartbreaker (Hbr): molecular marker in maize. *Proceeding of the National Academy of Science of the United State of America* **97**: 10083-10089.

- Chandler, M. and Mahillon, J. (2002). Insertion sequences revisited. In: Craig, N.L., Craigie, R., Gellert, M. and Lambowitz, A.M. (eds) *Mobile DNA II*. American Society for Microbiology Press. Washington. pp. 305-366.
- Chandrika, M. and Rai, V.R. (2009). Genetic fidelity in micropropagated plantlets of *Ochreinauclea missionis*: an endemic, threatened and medicinal tree using ISSR markers. *African Journal of Biotechnology* **8(13)**: 2933-2938.
- Chen, J.M., Stenson, P.D., Cooper, D.N. and Ferec, C. (2005). A systematic analysis of LINE-I endonuclease-dependent retrotranspositional events causing human genetic disease. *Human Genetics* **117**: 411-427.
- Chuang, S.J., Chen, C.L., Chen, J.J., Chou, W.Y. and Sung, J.M. (2009). Detection of somaclonal variation in micro-propagated *Echinacae purpurea* using AFLP markers. *Scientia Horticulturae* **120(1)**: 121-126.
- Collier, L.S. and Largaespada, D.A. (2007). Transposable elements and the dynamic somatic genome. *Genome Biology* **8(Suppl 1)**:S5 (DOI:10.1186/gb-2007-8-S1-S5): 1-5.
- Dam, A., Paul, S. and Bandyopadhyay, T.K. (2010). Direct somatic embryogenesis and plant regeneration from leaf explants of *Limonium sinensis* (Girard) Kuntze. *Scientia Horticulturae* **126**: 253-260.
- Debnath, M., Malik, C.P. and Bisen, P.S. (2006). Micropropagation: a tool for the production of high quality plant-based medicines. *Current Pharmaceutical Biotechnology* **7**: 33-49.
- d'Erfurth, I., Cosson, V., Eschstruth, A., Lucas, H., Kondorosi, A. and Ratet, P. (2003). Efficient transposition of the *Tnt1* tobacco retrotransposon in the model legume *Medicago truncatula*. *The Plant Journal* **34**: 95-106.
- De Setta, N., Costa, A.P.P., Lopes, F.R., Sluys, M.V. and Carareto, C.M.A. (2007). Transposon display supports transpositional activity of P elements in species of the *saltans* group of *Drosophila*. *Indian Academy of Science* **86(1)**: 37-43.
- Devos, K.M., Brown, J.K.M. and Bennetzen, J.L. (2002). Genome size reduction through illegitimate recombination counteracts genome expansion in *Arabidopsis*. *Genome Research* **12**: 1075-1079.
- Dobránski, J. and Teixeira de Silva, J.A. (2010). Micropropagation of apple- a review. *Biotechnology Advance* **28**: 462-488.
- Dooner, H.K. and Weil, C.F. (2007). Give-and-take: interaction between DNA transposon and their host plant genomes. *Current Opinion in Genetics and Development* **17(6)**: 486-492.
- Du, X.Y., Zhang, Q.L. and Luo, Z.R. (2009). Comparison of four molecular markers for genetics analysis in *Diospyros* L. (Ebenaceae). *Plant Systematics and Evolution* **281**: 171-181.
- Eka, C.R., Akomeah, P. and Asemota, O. (2005). Somatic embryogenesis in date palm (*Phoenix dactylifera* L.) from apical meristem tissues from 'zebia' and 'loko' landraces. *African Journal of Biotechnology* **4(3)**: 244-246.
- Elmeer, K.M.S., Gallagher, T.F. and Hennerty, M.J. (2009). RAPD-based detection of genomic instability in cucumber plants derived from somatic embryogenesis. *African Journal of Biotechnology* **8(14)**: 3219-3222.
- Eshraghi, P., Zarghami, R. and Ofoghi, H. (2005). Genetic stability of micropropagated plantlets in date palm. *Journal of Sciences, Islamic Republic of Iran* **16(4)**: 311-315.
- Evans, D.A. (1989). Somaclonal variation- genetic basis and breeding applications. *Trends in Genetics* **5(2)**: 46-50.
- Fay, M.F. (1992). Practical considerations in the development of a botanic garden micro-propagation laboratory. *Botanic gardens in changing worlds. The Proceedings of the Third International Botanic Gardens Conservation Congress. Rio de Janeiro.*
- Feschotte, C., Jiang, N. and Wessler, S.R. (2002). Plant transposable elements: Where genetics meets genomics. *Nature Review Genetics* **3**: 329-341.
- Fras, A. and Maluszynska, J. (2004). The correlation between the chromosome variation in callus and genotype of explants of *Arabidopsis thaliana*. *Genetica* **121**: 145-154.
- Fulneček, J. and Kovařík, A. (2014). How to interpret Methylation Sensitive Amplified Polymorphism (MSAP) profiles? *BMC Genetics* **15**: 2.
- Gatica-Arias, A.M., Arrieta-Espinoza, G. and Esquivel, A.M.E. (2008). Plant regeneration via indirect somatic embryogenesis and optimization of genetic transformation in *Coffea arabica* L. cvs. Caturra and Catuaí. *Electronic Journal of Biotechnology* **11(1)**: 1-12.
- Gaut, B.S. and Ross-Ibarra, J. (2008). Selection on major components of Angiosperm genomes. *Science* **320**: 484-486.
- Grandbastien, M.A. (1998). Activation of plant retrotransposons under stress conditions. *Trends in Plant Science* **3(5)**: 181-187.
- Hirochika, H. (1993). Activation of tobacco retrotransposons during tissue culture. *The EMBO Journal* **12**: 2521-2528.

- Hirochika, H., Sugimoto, K., Otsuki, Y., Tsugawa, H. and Kanda, M. (1996). Retrotransposon of rice involved in mutations induced by tissue culture. *Proceeding of National Academy of Science USA* **93**: 7783-7788.
- Huang, X., Lu, G., Zhao, Q., Liu, X. and Han, B. (2008). Genome-wide analysis of transposon insertion polymorphism reveals intraspecific variation in cultivated rice. *Genome Analysis* **148**: 25-40.
- Huang, J., Zhang, K., Shen, Y., Huang, Z., Li, M., Tang, D., Gu, M. and Cheng, Z. (2009). Identification of a high frequency transposon induced by tissue culture, *nDaiZ*, a member of the *hAT* family in rice. *Genomics* **93**: 274-281.
- Ikeda, R., Kokubu, C., Yusa, K., Keng, V.W., Horie, K. and Takeda, J. (2007). Sleeping Beauty transposase has an affinity for heterochromatin conformation. *Molecular Cell Biology* **27**: 1665-1676.
- Ito, H., Gaubert, H., Bucher, E., Mirouze, M., Vaillant, I. and Paszkowski, J. (2011). An siRNA pathway prevents transgenerational retrotransposition in plants subjected to stress. *Nature* **472**: 115-119.
- Ito, H. (2012). Small RNAs and transposon silencing in plants. *Development Growth and Differentiation* **54**: 100-107.
- Ito, H. (2013). Small RNAs and regulation of transposons in plants. *Genes and Genetic Systems* **88**: 3-7.
- Jaligot, E., Rival, A., Beule, T., Dussert, S. and Verdeil, J.L. (2000). Somaclonal variation in oil palm (*Elaeis guineensis* Jacq.): the DNA methylation hypothesis. *Plant Cell Reporter* **19**: 684-690.
- Jaligot, E., Beulé, T., Baurens, F.C., Billotte, N. and Rival, A. (2004). Search for methylation-sensitive amplification polymorphisms associated with the "mantled" variant phenotype in oil palm (*Elaeis guineensis* Jacq.). *Genome* **47**: 224-228.
- Jursch, T., Miskey, C., Izsvák, Z. and Ivics, Z. (2013). Regulation of DNA transposition by CpG methylation and chromatin structure in human cells. *Mobile DNA* **4**: 15.
- Kapitonov, V.V. and Jurka, J. (2008). A universal classification of eukaryotic transposable elements implemented in Repbase. *Nature Review Genetics* **9**: 411-412.
- Karajol, K. and Naik, G.R. (2011). Defective viral coat protein and retrotransposons might play role in NaCl stress tolerance in *Pigeon* pea (*Cajanus cajan*) cultivar. *World Journal of Science and Technology* **1(2)**: 17-25.
- Kazazian, H.H. Jr. (2004). Mobile elements: drivers of genome evolution. *Science* **303**: 1626-1632.
- Kaeppler, S.M., Kaeppler, H.F. and Rhee, Y. (2000). Epigenetic aspects of somaclonal variation in plants. *Plant Molecular Biology* **43**: 179-188.
- Kikuchi, K., Terauchi, K., Wada, M. and Hirano, H. (2003). The plant MITE mPing is mobilized in anther culture. *Nature* **421**: 167-170.
- Kohany, O., Gentles, A.I., Hankus, L. and Jurka, J. (2006). Annotation, submission & sequencing of repetitive elements in Repbase: Repbase Submitter and Censor. *BMC Bioinformatics* **7**: 474.
- Kolomietz, E., Meyn, M.S., Pandita, A. and Squire, J.A. (2002). The role of Alu repeat clusters as mediators of recurrent chromosomal aberrations in tumors. *Gene Chromosomes Cancer* **35**: 97-112.
- Krauss, V., Eisenhardt, C. and Unger, T. (2009). The genome of the stick insect *Medauroidea extradentata* is strongly methylated within genes and repetitive DNA. *PLoS ONE* **4(9)**: e7223. doi:10.1371/journal.pone.0007223.
- Krikorian, A.D. and Berquam, D.L. (1969). Plant cell and tissue cultures: the role of Haberlandt. *Botanical Review* **35**: 59-87.
- Kour, G., Kour, B., Kalu, S. and Dhar, M.K. (2009). Genetic and epigenetic instability of amplification-prone sequences of a novel B chromosome induced by tissue culture in *Plantago lagopus* L. *Plant Cell Report* **28**: 1857-1867.
- Kuan, L.I., Chuang, W., Shan, Y. and Hui-Xia, S. (2010). Comparing genetics characteristic of retrotransposon *TOS17* during different tissue culture processes in the rice cultivars Nipponbare and Shishoubaimao. *Agricultural Science in China* **9(2)**: 157-162.
- Kubis, S.E., Castilho, A.M.M.F., Vershinin, A.V. and Heslop-Harrison, J.S. (2003). Retroelement, transposon and methylation status in the genome of oil palm (*Elaeis guineensis*) and the relationship to somaclonal variation. *Plant Molecular Biology* **52**: 69-79.
- Le, Q.H. and Bureau, T. (2004). Prediction and quality assessment of transposon insertion display data. *BioTechniques* **36**: 222-228.
- Lenoir, A., Lavie, L., Prieto, J.L., Goubely, C., Côte, J.-C., Pelissier, T. and Deragon, J.M. (2001). The evolutionary origin and genomic organization of SINEs in *Arabidopsis thaliana*. *Molecular Biology and Evolution* **18(12)**: 2315-2322.
- Lisch, D. (2009). Epigenetic regulation of transposable elements in plants. *Annual Review of Plant Biology* **60**: 43-66.

- Lippman, Z. and Martienssen, R. (2004). The role of RNA interference in heterochromatic silencing. *Nature* **431**: 364-370.
- Lopes, T., Pinto, G., Loureiro, J., Costa, A. and Santos, C. (2006). Determination of genetic stability in long-term somatic embryogenic cultures and derived plantlets of cork oak using microsatellite markers. *Tree Physiology* **26**: 1145-1152.
- Lu, Y., Rong, T. and Cao, M. (2008). Analysis of DNA methylation in different maize tissues. *Journal of Genetics and Genomics* **35**: 41-48.
- Macas, J., Neumann, P. and Navrátilová, A. (2007). Repetitive DNA in the pea (*Pisum sativum* L.) genome: comprehensive characterization using 454 sequencing and comparison to soybean and *Medicago truncatula*. *BMC Genomics* **8**: 427.
- Mandal, P.K., Rawal, K., Ramaswamy, R., Bhattacharya, A. and Battacharya, S. (2006). Identification of insertion hot spots for non-LTR retrotransposons: computational and biochemical application to *Entamoeba histolytica*. *Nucleic Acid Research* **34(20)**: 5752-5763.
- Martienssen, R.A. and Colot, V. (2001). DNA methylation and epigenetic inheritance in plant and filamentous fungi. *Science* **293(5532)**: 1070-1074.
- Marum, L., Rocheta, M., Maroco, J., Oliveira, M.M. and Miguel, C. (2009). Analysis of genetic stability at SSR loci during somatic embryogenesis in maritime pine (*Pinus pinaster*). *Plant Cell Reports* **28**: 673-682.
- Matzke, M. and Birchler, J.A. (2005). RNAi-mediated pathways in the nucleus. *Nature Reviews Genetics* **6**: 24-35.
- Melayah, D., Bonnivard, E., Chalhoub, B., Audeon, C. and Grandbastien, M.A. (2001). The mobility of the tobacco retrotransposon correlates with its transcriptional activation by fungal factors. *The Plant Journal* **28**: 159-168.
- Meyers, B.C., Tingey, S.V. and Morgante, M. (2001). Abundance, distribution and transcriptional activity of repetitive elements in the maize genome. *Genome Research* **11**: 1660-1676.
- Michiba, K., Okamoto, T. and Mii, M. (2001). Increasing ploidy level in cell suspension culture of *Doritaenopsis* by exogenous application of 2,4-dichlorophenoxyacetic acid. *Plant Physiology* **112**: 142-148.
- Mirouze, M., Reinders, J., Bucher, E., Nishimura, T., Schneeberger, K., Ossowski, S., Cao, J., Weigel, D., Paszkowski, J., and Mathieu, O. (2009). Selective epigenetic control of retrotransposition in *Arabidopsis*. *Nature* **461**: 427-430.
- Miyao, A., Tanaka, K., Murata, K., Sawaki, H., Takeda, S., Abe, K., Shinozuka, Y., Onosato, K. and Hirochika, H. (2003). Target site specificity of the *Tos17* retrotransposon shows a preference for insertion within genes and against insertion in retrotransposon-rich regions of the genome. *Plant Cell* **15**: 1771-1780.
- Miyao, A., Nakagome, M., Ohnuma, T., Yamagata, H., Kanamori, H., Katayose, Y., Takahashi, A., Matsumoto, T. and Hirochika, H. (2012). Molecular spectrum of somaclonal variation in regenerated rice revealed by whole-genome sequencing. *Plant Cell Physiology* **53**: 256-264.
- Momose, M., Abe, Y. and Ozeki, Y. (2010). Miniature inverted-repeat transposable elements of *Stowaway* are active in potato. *Genetics* **186**: 59-66.
- Morgante, M., Hanafey, M. and Powell, W. (2002). Microsatellites are preferentially associated with non-repetitive DNA in plant genomes. *Nature Genetics* **30**: 194-200.
- Nontaswatsri, C. and Fukai, S. (2005). Regenerative callus of *Dianthus* 'Telstar Scarlet' showing mixoploidy produce diploid plants. *Plant Cell, Tissue and Organ Culture* **83**: 351-355.
- Ostertag, E.M. and Kazazian, H.H.Jr. (2001). Biology of mammalian retrotransposons. *Annual Review of Genetics* **35**: 501-538.
- Ostertag, E.M., Goodier, J.L., Zhang, Y. and Kazazian, H.H.Jr. (2003). SVA elements are non-autonomous retrotransposon that cause disease in human. *The American Journal of Human Genetics* **73**: 1444-1451.
- Ouyang, S. and Buell, C.R. (2004). The TIGR Plant Repeat Databases: a collective resource for the identification of repetitive sequences in plants. *Nucleic Acids Res.* 32(Database issue): D360-3.
- Paek, K.Y., Hanh, E.J. and Park, S.Y. (2011). Micropropagation of *Phalaenopsis* orchids via protocorms and protocorm-like body. *Methods in Molecular Biology* **710(4)**: 293-306.
- Parisod, C., Salmon, A., Zerjal, T., Tenaillon, M., Grandbastien, M.A. and Ainouche, M. (2009). Rapid structural and epigenetic reorganization near transposable elements in hybrid and allopolyploid genomes in *Spartina*. *New Phytologist* **184**: 1003-1015.
- Parisod, C., Salmon, A., Ainouche, M. and Grandbastien, M.A. (2014). Detecting epigenetic effects of transposable elements in plants. In: Charles Spillane and Peter C. McKeown (eds.). *Plant Epigenetics and Epigenomics: Methods and Protocols, Methods in Molecular Biology*, vol. 1112. Springer Science & Business Media, New York. pp. 211-217.

- Peraza-Echeverria, S., Herrera-Valencia, V.A. and James-Kay, A. (2001). Detection of DNA methylation changes in micropropagated banana plants using methylation-sensitive amplification polymorphism (MSAP). *Plant Science* **71**: 359-367.
- Picault, N., Chaparro, C., Piegu, B., Stenger, W., Formey, D., Llauro, C., Descombin, J., Sabot, F., Lasserre, E., Meynard, D., Guiderdoni, E. and Panaud, O. (2009). Identification of an active LTR retrotransposon in rice. *Plant Journal* **58**: 754-765.
- Piegu, B., Guvot, R., Picault, N., Roulin, A., Sanival, A., Kim, H., Cllura, K., Brar, D.S., Jackson, S., Wing, R.A. and Panaud, O. (2006). Doubling genome size without polyploidization: dynamics of retrotransposition-driven genomic expansions in *Oryza australiensis*, a wild relative of rice. *Genome Research* **16(10)**: 1262-1269.
- Pontaroli, A.C. and Camadro, E.L. (2005). Somaclonal variation in *Asparagus officinalis* plants regenerated by organogenesis from long-term callus cultures. *Genetic and Molecular Biology* **28(3)**: 423-430.
- Puente, R., González, A.I., Ruiz, M.L. and Polanco, C. (2008). Somaclonal variation in rye (*Secale cereals* L.) analyzed using polymorphic and sequenced AFLP markers. *In Vitro Cell & Development Biology-Plant* **44**: 419-426.
- Qin, Y., Li, H.L. and Guo, Y.D. (2007). High frequency embryogenesis, regeneration of broccoli (*Brassica oleracea* var. *italica*) and analysis of genetic stability by RAPD. *Scientia Horticulturae* **111**: 203-208.
- Rasheed, S., Fatima, T., Husnain, T., Bashir, K. and Riazuddin, S. (2005). RAPD characterization of somaclonal variation in *indica* basmati rice. *Pakistan Journal of Botany* **37(2)**: 249-262.
- Reyna-Lopez, G.E., Simpson, J. and Ruiz-Herrera, J. (1997). Differences in DNA methylation patterns are detectable during the dimorphic transition of fungi by amplification of restriction polymorphisms. *Molecular Genomics and Genetics* **253(6)**: 703-710.
- Rout, G.R., Mohapatra, A. and Mohan Jain, S. (2006). Tissue culture of ornamental pot plant: A critical review on present scenario and future prospect. *Biotechnology Advance* **24**: 531-560.
- Rozen, S. and Skaletsky, H.J. (2000). Primer3 on the WWW for general users and for biologist programmers. In: Krawetz, S. and Misener, S. (eds) *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press. New Jersey. pp. 365-386.
- Sabot, F., Picault, N., El-Baidouri, M., Llauro, C., Chaparro, C., Piegu, B., Roulin, A., Guiderdoni, E., Delabastide, M., McCombie, R. and Panaud, O. (2011). Transpositional landscape of the rice genome revealed by paired-end mapping of high-throughput re-sequencing data. *Plant Journal* **66**: 241-246.
- Sen, S.K., Han, K., Wang, J., Lee, J., Wang, H., Callinan, P.A., Dyer, M., Cordaux, R., Liang, P. and Batzer, M.A. (2006). Human genomic deletions mediated by recombinant between Alu elements. *The American Journal of Human Genetics* **79**: 41-53.
- Shirasu, K., Schulman, A.H., Lahaye, T. and Schulze-Lefert, P. (2000). A contiguous 66 kb barley DNA sequence provides evidence for reversible genome expansion. *Genome Research* **10**: 908-915.
- Sianipar, F.S., Wattimena, G.A., Soehartono, M.T., Aswidinnoor, H., Toruan Mathius, N. and Ginting, G. (2008). DNA methylation detection of oil palm (*Elaeis guineensis* Jacq.) somatic embryo by randomly amplified DNA fingerprinting with methylation-sensitive (RAF-SM) and RP-HPLC. *Journal of Biotechnology Research in Tropical Region* **1**: 1-6.
- Singh, M.P. and Kumar, S. (2009). *Principal and methodology in plant tissue culture*. APH Publishing, India. pp. 9-56.
- Slotkin, R.K. and Martienssen, R. (2007). Transposable elements and the epigenetic regulation of the genome. *Nature Reviews Genetics* **8**: 272-285.
- Soniya, E.V., Banerjee, N.S. and Das, M.R. (2001). Genetic analysis of somaclonal variation among callus-derived plants of tomato. *Current Science* **80(9)**: 1213-1215.
- Syed, N.H. and Flavell, A.J. (2007). Sequence-specific amplification polymorphisms (SSAPs): a multi-locus approach for analyzing transposon insertions. *Nature Protocol* **6(1)**: 2746-2752.
- Tahara, M., Aoki, T., Suzuka, S., Yamashita, H., Tanaka, M., Matsunaga, S. and Kokumai, S. (2004). Isolation of an active element from a high-copy-number family of retrotransposons in the sweetpotato genome. *Molecular Genetics and Genomics* **272**: 116-127.
- Takata, M., Kiyohara, A., Takasu, A., Kishima, Y., Ohtsubo, H. and Sano, Y. (2007). Rice transposable elements are characterized by various methylation environments in the genome. *BMC Genomics* **8**: 469.
- Takebe, I., Labib, G. and Melchers, G. (1971). Regeneration of whole plants from isolated mesophyll protoplasts of tobacco. *Naturwissenschaften* **58**: 318-320.

- Tanurdzic, M., Vaughn, M.W., Jiang, H., Lee, T.J., Slotkin, R.K., Sosinski, B., Thompson, W.F., Doerge, R.W. and Martienssen, R.A. (2008). Epigenomic consequences of immortalized plant cell suspension culture. *PLoS Biol* **6**: 2880-2895.
- Temel, A., Kartal, G. and Gozukirmizi, N. (2008). Genetic and epigenetic variations in barley calli culture. *Biotechnology and Biotechnological Equipment* **22(4)**: 911-914.
- Thorpe, T.A. (2007). History of plant tissue culture. *Molecular Biotechnology* **37**: 169-180.
- Todorovska, E. (2007). Retrotransposons and their role in plant-genome evolution. *Biotechnology and Biotechnological Equipment* **21(3)**: 294-305.
- Tsaftaris, A.S., Polidoros, A.N., Koumproglou, R., Tani, E., Kovacevic, N. and Abatzidou, E. (2005). Epigenetic mechanism in plants and their implications in plant breeding. In: Tuberosa, R., Phillips, R.L. and Gale, M. (eds). *Proceedings of the International Congress "In the Wake of the Double Helix: From the Green Revolution to the Gene Revolution"*. Avenue Media. Italy. pp. 157-171.
- Vasil, V. and Hildebrand, A.C. (1965). Differentiation of tobacco plants from single, isolated cells in micro-cultures. *Science* **150**: 889-892.
- Vázquez, A.M. and Linacero, R. (2010). Stress and somaclonal variation. In: Pua E.-C., Davey M. (eds). *Plant Developmental Biology – Biotechnological Perspectives*, vol. 2. Springer, New York. pp 45–64.
- Venkatachalam, L. and Sreedhar, R.V. (2007). Micropropagation in banana using high levels of cytokinins does not involve any genetic changes as revealed by RAPD and ISSR markers. *Plant Growth Regulation* **51**: 193-205.
- Venturi, S., Dondini, L., Donini, P. and Sansavini, S. (2006). Retrotransposon characterization and fingerprinting of apple clones by S-SAP markers. *Theoretical and Applied Genetics* **112(3)**: 440-444.
- Vergara-Galicia, J., Aguirre-Crespo, F., Castillo-España, P., Arroyo-Mora, A., López-Escamilla, A.L., Villalobos-Molina, R. and Estrada-Soto, S. (2010). Micropropagation and vasorelaxant activity of *Laelia autumnalis* (Orchidaceae). *Natural Product Research* **24(2)**: 106-114.
- Vizel, M., Loya, Y., Downs, C.A. and Kramarsky-Winter, E. (2010). A novel method for coral explants culture and micropropagation. *Marine Biotechnology* (DOI 10.1007/s10126-010-9313-z): 1-10.
- Wegscheider, E., Benjak, A. and Forneck, A. (2009). Clonal variation in Pinot noir revealed by S-SAP involving universal retrotransposon-based sequences. *American Journal of Enology and Viticulture* **60(1)**: 104-109.
- Wicker, T., Sabot, F., Hua-Van, A., Bennetzen, J.L., Capy, P., Chalhoub, B., Flavell, A., Leroy, P., Morgante, M., Panaud, O., Paux, E., San Miguel, P. and Schulman, A.H. (2007). A unified classification system for eukaryotic transposable elements. *Natural Review Genetics* **8**: 973-982.
- Xiong, L.Z., Xu, C.G., Saghai, M.M.A. and Zhang, Q. (1999). Patterns of cytosine methylation in an elite rice hybrid and its parental lines, detected by a methylation-sensitive amplification polymorphism technique. *Molecular and General Genetics* **261**: 439-446.
- Yusa, K., Takeda, J. and Horie, K. (2004). Enhancement of Sleeping Beauty transposition by CpG methylation: possible role of heterochromatin formation. *Molecular Cell Biology* **24**: 4004-4018.
- Zedek, F., Šmerda, J., Šmarda, P. and Bureš, P. (2010). Correlated evolution of LTR retrotransposons and genome size in the genus *Eleocharis*. *BMC Plant Biology* **10**: 265.
- Zerjal, T., Joets, J., Alix, K., Grandbastien, M.A. and Tenailon, M.I. (2009). Contrasting evolutionary patterns and target specificities among three *Tourist*-like MITE families in the maize genome. *Plant Molecular Biology* **71**: 99-114.